

3-Dimensional Lattice Polytopes Without Interior Lattice Points

Jaron Treutlein

ABSTRACT. A theorem of Howe states that every 3-dimensional lattice polytope P whose only lattice points are its vertices, is a Cayley polytope, i.e. P is the convex hull of two lattice polygons with distance one.

We want to generalize this result by classifying 3-dimensional lattice polytopes without interior lattice points. The main result will be, that they are up to finite many exceptions either Cayley polytopes or there is a projection, which maps the polytope to the double unimodular 2-simplex.

To every such polytope we associate a smooth projective surface of genus 0.

1. Introduction

Let $M \cong \mathbb{Z}^n$ be a lattice and $P \subset M \otimes_{\mathbb{Z}} \mathbb{R}$ be an n -dimensional lattice polytope, i.e. the set of vertices of P is contained in M . However in the following we will always suppose $M = \mathbb{Z}^n$.

Let $P \subset \mathbb{R}^3$ be a 3-dimensional lattice polytope. If $|P^\circ \cap \mathbb{Z}^3| = i > 0$, then we will derive from Hensley's theorem [5], that its volume $\text{vol}(P)$ is bounded by a constant depending only on i . Jeffrey Lagarias and Günter M. Ziegler proved in [9]:

THEOREM 1.1 (Lagarias, Ziegler). *Let F be a family of n -dimensional lattice polytopes. Then is equivalent:*

- (1) $\text{vol}(P) < C \ \forall P \in F$ with a constant $C > 0$.
- (2) Up to affine unimodular transformation, F is a finite set.

Thus the finiteness up to affine unimodular transformation of 3-dimensional lattice polytopes having exactly $i > 0$ interior lattice points follows. So, it remains to consider the case $i = 0$ in order to classify all 3-dimensional lattice polytopes.

Let $\{e_1, \dots, e_n\} \subset \mathbb{R}^n$ be the standard basis of $\mathbb{R}^n$. We denote the unimodular n -simplex $\text{conv}(0, e_1, \dots, e_n)$ by Δ_n . If two lattice polytopes P and Q are equivalent modulo affine unimodular transformation, we will describe it by $P \cong Q$.

John R. Arkinstall [1], Askold Khovanskii [7], Robert J. Koelman [8] and Josef Schicho [14] proved that a lattice polygon without interior lattice points satisfies $P \cong 2\Delta_2$ or $P \cong \text{conv}(0, e_1, h_1 e_2, e_1 + h_2 e_2)$ with some heights $h_1, h_2 \in \mathbb{N}$. The latter is called a Lawrence polygon. Thus a lattice polygon without interior lattice

1991 *Mathematics Subject Classification.* Primary 52B20, 14M25; Secondary 14Q10.

Key words and phrases. Lattice polytopes, Howe, Surfaces.

points either has a projection to Δ_1 , which is the only 1-dimensional lattice polytope without interior lattice points, or it is equivalent to $2\Delta_2$.

Victor Batyrev conjectured that any 3-dimensional lattice polytope without interior lattice points either can be projected to Δ_1 , to $2\Delta_2$ or its volume is bounded. From Theorem 1.1 it will follow that up to unimodular transformation there is only a finite number of exceptional polytopes of this last class.

A lattice polytope $P \subset \mathbb{R}^n$ is described as the Cayley polytope of the lattice polytopes $P_0, \dots, P_{r-1} \subset \langle e_r, \dots, e_n \rangle$, $1 < r \leq n$, if $P \cong \text{conv}(0 \times P_0, e_1 \times P_1, \dots, e_{r-1} \times P_{r-1})$. Then we call P simply a Cayley polytope and notice that these are the lattice polytopes which can be projected to Δ_1 .

An n -dimensional lattice polytope is called a k -fold lattice pyramid over an $(n-k)$ -dimensional lattice polytope Q , if it is the Cayley polytope of Q and k lattice points.

It is clear, that lattice polytopes having a projection to Δ_1 or $2\Delta_2$, both have no interior lattice points.

Roger Howe proved in 1977 the following:

THEOREM 1.2 (Howe). *Let $P \subset \mathbb{R}^3$ be a lattice polytope whose only lattice points are vertices. Then $|P \cap \mathbb{Z}^3| \leq 8$ and P is a Cayley polytope.*

As he did not publish it, Herbert E. Scarf did it [13]. There are further proofs for example by G.K. White [15], Bruce Reznick [11], András Sebő [12], David R. Morrison and Glenn Stevens [10].

The aim of this paper is to prove Victor Batyrev's conjecture and hence generalize Theorem 1.2:

THEOREM 1.3. *Let $P \subset \mathbb{R}^3$ be a lattice polytope without interior lattice points. Then either is P a Cayley polytope, P can be projected to the double unimodular 2-simplex or P is an exceptional polytope, whereas up to unimodular transformation there is only a finite number of these.*

Some of these exceptional simplices ($P_1, \dots, P_6$) are in figure 1. All the others (there are 15 more simplices) are included in one of them. Thus we describe $P_1, \dots, P_6$ as maximal exceptional lattice simplices.

Moreover there are maximal exceptional lattice polytopes which are not simplices, see for example P_7, P_8 and P_9 in figure 2. It is not known if these are the only maximal lattice polytopes.

These are the known maximal exceptional lattice polytopes:

$$\begin{aligned}
P_1 &:= \text{conv}(0, e_1, 2e_1 + 5e_2, 3e_1 + 5e_3) \\
P_2 &:= \text{conv}(0, 3e_1, e_1 + 3e_2, 2e_1 + 3e_3) \\
P_3 &:= \text{conv}(0, 3e_1, 3e_2, 3e_3) \\
P_4 &:= \text{conv}(0, 4e_1, 4e_2, 2e_3) \\
P_5 &:= \text{conv}(0, 4e_1, 2e_1 + 4e_2, e_1 + 2e_3) \\
P_6 &:= \text{conv}(0, 6e_1, 3e_2, 2e_3) \\
P_7 &:= \text{conv}(\pm 2e_1, \pm 2e_2, e_1 + e_2 + 2e_3) \\
P_8 &:= \text{conv}(\pm e_1, 2e_2, e_1 + 2e_3 \pm e_1, e_1 + 2e_2 + 2e_3) \\
P_9 &:= \text{conv}(\pm e_1, \pm e_2, e_1 + e_2 + 2e_3 \pm e_1, e_1 + e_2 + 2e_3 \pm e_2).
\end{aligned}$$




FIGURE 1. The maximal exceptional lattice simplices




FIGURE 2. Some more maximal exceptional lattice polytopes

CONJECTURE 1.4. Up to a finite number of exceptional lattice polytopes to every n -dimensional lattice polytope without interior lattice points, there is a projection mapping it onto an m -dimensional lattice polytope without interior lattice points, $m < n$.

EXAMPLE 1.5. Christian Haase and Günter M. Ziegler showed in [4] that the only lattice points of the 4-dimensional lattice simplex $\text{conv}(e_1, e_2, e_3, e_4, 2e_1 + 2e_2 + 3e_3 + (k - 6)e_4)$ are its vertices, if and only if $\gcd(k, 6) = 1$. Moreover they say that the only lattice points of the 5-dimensional lattice simplex $\text{conv}(0, e_1, e_2, e_3, e_1 + e_2 + e_3 + 6e_4, 2e_1 + 3e_2 + 4e_3 + 9e_5)$ are its vertices. It is easy to see that there are projections mapping them onto $2\Delta_2$.

Now we will show how you can get a smooth projective surface associated to a 3-dimensional lattice polytope:

Given a lattice polytope $P \subset \mathbb{R}^3$, consider the normal fan $\Sigma(P)$ defined by the cones

$$\sigma(\theta) := \{y \in (\mathbb{Z}^3)^* : \min_{x \in P} \langle x, y \rangle = \langle z, y \rangle \ \forall z \in \theta\},$$

whereas θ is a face of P .

Denote by $\mathbb{P}_P$ the projective toric variety defined by this fan over a field K . Now consider the compactification $\mathcal{S}_P \subset \mathbb{P}_P$ of the surface

$$\left\{ \sum_{v \in P \cap \mathbb{Z}^3} a_v X_1^{v_1} X_2^{v_2} X_3^{v_3} = 0 \right\} \subset (\overline{K}^*)^3 \hookrightarrow \mathbb{P}_P$$

in $\mathbb{P}_P$, whereas $\overline{K}$ denotes the algebraic closure of K and the coefficients $a_v \in K$ satisfy the following generic condition:

$$\left\{ \sum_{v \in \theta \cap \mathbb{Z}^3} a_v X_1^{v_1} X_2^{v_2} X_3^{v_3} = 0 \right\} \subset (\overline{K}^*)^3$$

is smooth for every face $\theta \subset P$.

A.G. Khovanskii showed in [6] that the number of interior lattice points of $P \subset \mathbb{R}^3$ is the same as the geometric genus of the surface $\mathcal{S}_P$. So by considering lattice polytopes $P \subset \mathbb{R}^3$ without interior lattice points we get – after resolution of singularities – some smooth projective surfaces of genus 0, i.e. smooth projective surfaces which are “nearly” rational.

In fact if P is a Cayley polytope of two polygons, the corresponding surface will be rational. However if P has a projection onto $2\Delta_2$ we will get a conic bundle, which – in general – will be not rational. Now consider the maximal exceptional lattice polytopes.

$\mathcal{S}_{P_1}$ is the Godeaux surface

$$\left\{ \sum_{i=1}^4 x_i^5 = 0, \ x \in \mathbb{P}^3 \right\} / \mu_5,$$

whereas $\zeta.x_i := \zeta^i x_i$, $\zeta^5 = 1$.

$\mathcal{S}_{P_5}, \mathcal{S}_{P_7}, \mathcal{S}_{P_8}$ and $\mathcal{S}_{P_9}$ are Enriques surfaces. $\mathcal{S}_{P_5}$ can be realized as

$$\mathcal{S}_{P_5} := \{x_1^8 + x_2^8 + x_3^4 + x_4^2 = 0, \ x \in \mathbb{P}^3(1, 1, 2, 4)\} / \mu_2,$$

whereas $(-1).x := (-x_1 : x_2 : x_3 : x_4)$.

$\mathcal{S}_{P_8}$ is the closure of

$$\{(u : v), (x : y : z) \in \mathbb{P}^1 \times \mathbb{P}^2(1, 1, 2) : (u^2 + v^2)(x^4 y^4 + x^4 z^2 + y^4 z^2) = 0\} / \mu_2,$$

whereas $(-1).(u : v), (x : y : z) := ((u : -v), (x : -y : -z))$.

$\mathcal{S}_{P_9}$ is the closure of

$$\left\{ ((x_0 : x_1), (y_0 : y_1), (z_0 : z_1)) \in \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 : \right. \\ \left. (x_0^2 + x_1^2)(y_0^2 + y_1^2)(z_0^2 + z_1^2) = 0 \right\} / \mu_2,$$

whereas $(-1).(x, y, z) := ((x_0 : -x_1), (y_0 : -y_1), (z_0 : -z_1))$.

$\mathcal{S}_{P_3}$ is the cubic del Pezzo surface

$$\left\{ \sum_{i=1}^4 x_i^3 = 0, \ x \in \mathbb{P}^3 \right\}.$$

$\mathcal{S}_{P_2}$ is the quasi-homogeneous nonic in weighted projective space

$$\mathcal{S}_{P_2} := \{x_1^9 + x_2^9 + x_3^9 + x_4^3 = 0, x \in \mathbb{P}^3(1, 1, 1, 3)\} / \mu_9,$$

whereas $\zeta.x := (\zeta x_1 : \zeta^4 x_2 : x_3 : \zeta^6 x_4)$, $\zeta^9 = 1$.

$\mathcal{S}_{P_4}$ is the quasi-homogeneous quartic in weighted projective space

$$\{x_1^4 + x_2^4 + x_3^4 + x_4^2 = 0, x \in \mathbb{P}^3(1, 1, 1, 2)\}.$$

$\mathcal{S}_{P_6}$ is the quasi-homogeneous sextic in weighted projective space

$$\{x_1^6 + x_2^6 + x_3^3 + x_4^2 = 0, x \in \mathbb{P}^3(1, 1, 2, 3)\}.$$

The degree of a polytope $P \subset \mathbb{R}^n$ is the biggest integer $d \in \mathbb{N}$ such that

$$\left| \left((n+1-d)P \right)^\circ \cap \mathbb{Z}^n \right| > 0.$$

Consequently, the degree of a lattice polytope $P \subset \mathbb{R}^3$ without interior lattice points is 0, 1 or 2. The lattice polytopes of degree smaller than 2 are already classified (cf. [2]), which generalizes the result of Arkinstall, Khovanskii, Koelman and Schicho:

THEOREM 1.6 (Batyrev, Nill). *Let P be an n -dimensional lattice polytope. If $\deg(P) = 0$, then $P \cong \Delta_n$. If $\deg(P) = 1$, then P is an $(n-2)$ -fold lattice pyramid over $2\Delta_2$ or a Lawrence polytope, i.e. a Cayley polytope of n intervals.*

COROLLARY 1.7. Let $P \subset \mathbb{R}^n$, $n \geq 3$, be a lattice polytope and $\deg(P) \leq 1$. Then is P a Cayley polytope.

Let $P \subset \mathbb{R}^3$ be a lattice polytope without interior lattice points. Then $\deg(P) \leq 2$. Thus it remains to consider the case $\deg(P) = 2$.

In the following we will generalize Theorem 1.2 step by step in order to prove Theorem 1.3 in the end.

Acknowledgments: The author would like to thank Victor Batyrev for discussions and joint work on this subject.

2. White Lattice Simplices

In the following we denote by $\text{Vol}(P) = n! \text{vol}(P)$ the normalized volume of an n -dimensional lattice polytope.

If $F \subset \mathbb{R}^2$ is a lattice polygon without interior lattice points, we derive $F \cong 2\Delta_2$ or F is a Lawrence polytope from section 1.

We describe a 3-dimensional lattice polytope as white, if all its lattice points are on its edges. In [15] G.K. White also proved the following generalization of Theorem 1.2, using inequalities of Gaussian brackets. We give a new proof:

PROPOSITION 2.1. Let $P \subset \mathbb{R}^3$ be a white lattice simplex with $F \not\cong 2\Delta_2$ for every facet $F \subset P$ of P . Then is P a Cayley polytope.

PROOF. As every facet of P is a Lawrence polygon, there are at most two edges of P of length bigger than 1, i.e. edges with more than two lattice points. By Theorem 1.2 we may assume that there is at least one edge of length bigger than 1.

Case 1 **There is exactly one edge of length $l_1 > 1$.**

Denote the vertices of P by $a, b, c, d \in \mathbb{Z}^3$ and let $f \in \text{conv}(a, b)^\circ \cap \mathbb{Z}^3$ be an interior point of the edge $\text{conv}(a, b)$.

Case A $l_1 = 2$

The lattice simplices $\Delta := \text{conv}(a, c, d, f)$ and $\Delta' := \text{conv}(b, c, d, f)$ are Cayley polytopes by Theorem 1.2. So Δ is stretched between parallel planes E_1, E_2 having distance 1, and Δ' is stretched between E'_1, E'_2 having also distance 1.

If a and f are both contained in E_1 or E_2 , then it will follow by linearity that b is also contained in the same plane. Hence P is a Cayley polytope stretched between E_1 and E_2 . The same argument holds if b and f are both contained in E'_1 or E'_2 .

If Δ is a pyramid, then $\Delta \cong \Delta_3$ will follow. So $P = \Delta \cup \Delta' \cong \text{conv}(-e_1, e_1, e_2, e_3)$ is a Cayley polytope. The same argument holds for Δ' . So assume in the following that Δ and Δ' are not pyramids.

Let $f = 0, c = e_1, d = e_2, b = -a$ and $0 \leq a_1, a_2 < a_3$. Notice that $\text{Vol}(\Delta) = \text{Vol}(\Delta') = a_3$. We may assume that Δ is a Cayley polytope of $\Delta_1 = \text{conv}(0, e_1)$ and $\Delta_2 = \text{conv}(e_2, a)$, i.e. $\text{Vol}(\text{conv}(0, e_1, a - e_2)) = \text{Vol}(\Delta) = a_3$. Hence $\gcd(a_2 - 1, a_3) = a_3$, which implies $a_3 | (a_2 - 1)$. So $a_2 = 1$. It remains to consider the following two cases:

Case a: Δ' is the Cayley polytope of $\Delta'_1 = \text{conv}(0, e_1)$ and $\Delta'_2 = \text{conv}(e_2, -a)$. Here is $\text{Vol}(\text{conv}(0, e_1, a + e_2)) = \text{Vol}(\Delta') = a_3$. Hence $\gcd(a_2 + 1, a_3) = a_3$, which implies $a_3 | (a_2 + 1) = 2$. As $a_3 > a_2 = 1$ we derive $a_3 = 2$. Then $a_1 \in \{0, 1\}$.
If $a_1 = 0$, then $P \subset \{0 \leq x_1 \leq 1\}$ will be a Cayley polytope.
If $a_1 = 1$, then e_1 and e_2 will have distance 1 to the plane $\text{lin}(a, e_1 - e_2)$, which includes $0, a, -a$. So again P will be a Cayley polytope.

Case b: Δ' is the Cayley polytope of $\Delta'_1 = \text{conv}(0, e_2)$ and $\Delta'_2 = \text{conv}(e_1, -a)$. Here is $\text{Vol}(\text{conv}(0, e_2, a + e_1)) = \text{Vol}(\Delta') = a_3$. Hence $\gcd(a_1 + 1, a_3) = a_3$, which implies $a_3 | (a_1 + 1)$. Then e_1 and e_2 have distance 1 to the plane $\text{lin}(a, e_1 - e_2)$, which includes $0, a, -a$. Thus P is a Cayley polytope.

Case B $l_1 > 2$

By induction Δ and Δ' are both Cayley polytopes, whereas we adopt the same notation as before. Without loss of generality let $|\text{conv}(a, f)^\circ \cap \mathbb{Z}^3| > 0$. Then a and f – and thus also b – are contained in E_1 or E_2 . Hence, P is a Cayley polytope stretched between E_1 and E_2 .

Case 2 **There are two edges of lengths $l_1 > 1, l_2 > 1$.**

Let $l_1 + 1 = |\text{conv}(a, b) \cap \mathbb{Z}^3|$ and $l_2 + 1 = |\text{conv}(c, d) \cap \mathbb{Z}^3|$. There is a lattice point $f \in \text{conv}(a, b)^\circ \cap \mathbb{Z}^3$ such that $|\text{conv}(b, f)^\circ \cap \mathbb{Z}^3| = 0$.

By induction, $\Delta^{(1)} := \text{conv}(a, c, d, f)$ and $\Delta^{(2)} := \text{conv}(b, c, d, f)$ are Cayley polytopes, i.e. there is a number $z^{(i)} \in \mathbb{Z}$ and a linear functional

$y^{(i)} \in (\mathbb{Z}^3)^*$ in such a way, that $\Delta^{(i)}$ is contained in $W^{(i)} := \{x \in \mathbb{R}^3 : z^{(i)} \leq \langle x, y^{(i)} \rangle \leq z^{(i)} + 1\}$, $i \in \{1, 2\}$.

The lattice polygon $\text{conv}(c, d, f)$ is a Lawrence polygon. So every facet of $\Delta^{(i)}$, $i \in \{1, 2\}$ is a Lawrence polygon.

If a and f are both contained in the same boundary plane of $W^{(1)}$ then b will be in it too. In this case $P \subset W^{(1)}$ is a Cayley polytope. So let a and f be in different boundary planes of $W^{(1)}$. Thus $l_1 = 2$. As $l_2 > 1$ we notice that c and d are in the same boundary plane of $W^{(i)}$.

Hence $\Delta^{(i)}$ is a pyramid for $i \in \{1, 2\}$ and thus a Lawrence polytope. We may choose $\Delta^{(1)} = \text{conv}(0, e_1, l_2 e_2, e_1 + e_3)$. So we see that $P \subset \{0 \leq X_1 \leq 1\}$ is a Cayley polytope. $\square$

PROPOSITION 2.2. Let $P \subset \mathbb{R}^3$ be a white lattice simplex. Then is P a Cayley polytope or $P \cong 2\Delta_3$.

PROOF. Because of Proposition 2.1 we may assume $P = \text{conv}(0, 2e_1, 2e_2, d)$ with $d \in \mathbb{Z}^3$, $d_3 > 1$.

If there is a facet $F \cong 2\Delta_2$ of P , $F \neq P \cap X_3^\perp$, then every facet of P will be of this kind and $\frac{1}{2}P$ will be a lattice polytope of degree 0 or 1 having only unimodular facets. So $P \cong 2\Delta_3$.

Now let every facet $F \neq P \cap X_3^\perp$ be a Lawrence polygon. As P is a simplex and $v := (v_1, v_2, 1) \notin P \forall v_1, v_2 \in \mathbb{Z}$, the line g through d and v intersects with P only in d . In particular $g \cap X_3^\perp \not\subset 2\Delta_2$. Consequently $M \cap 2\Delta_2 = \emptyset$, whereas

$$M = \left\{ v_1 \cdot \frac{d_3}{d_3 - 1} - w_1, v_2 \cdot \frac{d_3}{d_3 - 1} - w_2 : v_1, v_2 \in \mathbb{Z} \right\}$$

with some $w_1, w_2 \in \mathbb{Z}$. But this is a contradiction to the fact that $\emptyset \neq M \cap \{0 \leq X_1, X_2 \leq 1\} \subset 2\Delta_2$. So, $d_3 = 1$ and P is a Cayley polytope. $\square$

3. White Lattice Polytopes

PROPOSITION 3.1. Let F be a lattice polygon, $d \in \mathbb{Z}^3$ and $P = \text{conv}(F, d) \subset \mathbb{R}^3$ be a white lattice polytope. Then is P a Cayley polytope.

PROOF. As $|F^\circ \cap \mathbb{Z}^3| = 0$, we may suppose F is a Lawrence polytope but not a simplex by Theorem 1.6 and Proposition 2.2. So let F be the Cayley polytope of two parallel edges $f^{(1)}$ and $f^{(2)}$. Assume that $X_3(f^{(1)}) = X_3(f^{(2)}) = 0$ and $X_1(f^{(1)}) = 0, X_1(f^{(2)}) = 1$.

Choose two intervals $\Delta_1 \cong q^{(i)} \subset f^{(i)}$, $i \in \{1, 2\}$, and consider the lattice subpolytope $Q := \text{conv}(q^{(1)}, q^{(2)}, d) \subset P$. Let $0 \leq d_1, d_2 < d_3$ and suppose $d_3 > 1$.

Assume that the only lattice points of Q are its vertices. Then is Q a Cayley polytope by Theorem 1.2. But as all the facets of Q have no interior lattice points we conclude that every subsimplex is unimodular and hence $d_3 = 1$, which is a contradiction. So we may assume that $\text{conv}(0, d)^\circ \neq \emptyset$, $q_1 = \text{conv}(0, e_2)$ and $q_2 = \text{conv}(e_1, e_1 + e_2)$. We will show that $d_1 = 0$, which will imply that $P \subset \{0 \leq X_1 \leq 1\}$.

Subdivide Q into $\Delta^{(1)} := \text{conv}(0, e_1, e_1 + e_2, d)$ and $\Delta^{(2)} := \text{conv}(0, e_2, e_1 + e_2, d)$. By Proposition 2.2 they are both Cayley polytopes of normalized volume d_3 .

If $\Delta^{(1)}$ is a pyramid, then it will be a Lawrence simplex and hence $d_1 = d_2 = 0$.

So let $\Delta^{(1)}$ be a Cayley polytope of $\text{conv}(0, d)$ and $\text{conv}(e_1, e_1 + e_2)$ resp. $\text{conv}(e_2, e_1 + e_2)$. Then is $\text{Vol}(\text{conv}(0, e_2, d)) = \text{Vol}(\Delta) = d_3$. Thus $\gcd(d_1, d_3) = d_3$, which implies $d_3 | d_1$. Hence $d_1 = 0$. $\square$

PROPOSITION 3.2. Let $P \subset \mathbb{R}^3$ be a white lattice polytope with at most 5 vertices. Then is P a Cayley polytope or $P \hookrightarrow 3\Delta_3$ and $(2\Delta_2)^\circ \hookrightarrow P^\circ$.

PROOF. By the Propositions 2.2 and 3.1 we may assume that P is a circuit, i.e. the convex hull of a 2-dimensional lattice simplex Δ and a 1-dimensional lattice polytope $\text{conv}(a, b)$, each without interior lattice points, such that $\text{conv}(a, b)^\circ, \Delta^\circ \subset P^\circ$. By Theorem 1.2 we may moreover assume that there is an edge of P of length $n \geq 2$. Consider the following two cases:

Case A: $\Delta \cong \Delta_2$.

There are lattice points $c, d \in P \cap \mathbb{Z}^3$ such that $\text{conv}(\Delta, c, d)$ is a circuit whose only lattice points are its vertices. By Theorem 1.2 we may suppose that $\text{conv}(\Delta, c, d) = \text{conv}(0, e_1, e_2, -e_3, f)$ with $\Delta = \text{conv}(0, e_1, e_2)$, $f = e_1 + xe_2 + ye_3$ and $x, y \in \mathbb{N}$. As P is a circuit we see moreover $x > 0$ and $1 + x < y$, in particular $y > 1$. In order to get P back, it remains to elongate f or $-e_3$ from Δ . If we elongate f from e_1 , then $P \subset \{0 \leq X_1 \leq 1\}$ will be a Cayley polytope. The same will be if we elongate $-e_3$ from 0 or from e_2 . We will show now by assuming the contrary that in the remaining three cases you cannot reach any subpolytope of P :

Case 1: Elongate f from 0.

Here we get the circuit $\text{conv}(0, e_1, e_2, -e_3, 2f) \subset P$. By Proposition 2.2 is the subsimplex $\text{conv}(0, e_1, e_2, 2f)$ a Cayley polytope.

If it was a pyramid, then it would follow from Theorem 1.6 that $f_3 | f_1$ and $f_3 | f_2$, which would be a contradiction.

Thus $\text{conv}(0, e_1, e_2, 2f)$ is the Cayley polytope of $\text{conv}(0, 2f)$ and $\text{conv}(e_1, e_2)$. So $\text{Vol}(\text{conv}(0, 2f, e_1 - e_2)) = \text{Vol}(\text{conv}(0, e_1, e_2, 2f)) = 2y$ and consequently $2x + 2 = 2y$, which is a contradiction to the fact that $1 + x < y$.

Case 2: Elongate f from e_2 .

Here we get the circuit $\text{conv}(0, e_1, e_2, -e_3, 2f - e_2) \subset P$. As before we receive $2y = \text{Vol}(\text{conv}(0, 2f - e_2, e_1 - e_2))$, which implies $2y | (2x + 1)$, which is a contradiction.

Case 3: Elongate $-e_3$ from e_1 .

Here we get the circuit $\text{conv}(0, e_1, e_2, -e_1 - 2e_3, f)$. So $y = 1$, which is again a contradiction.

Case B: There is an edge of Δ of length $n \geq 2$.

By Theorem 1.6 we may assume $\Delta = \text{conv}(0, ne_1, \delta e_2)$ with $\delta \in \{1, 2\}$, $a_3 > 0 > b_3$. Subdivide P into $\Delta' := P \cap \{X_3 \geq 0\}$ and $\Delta'' := P \cap \{X_3 \leq 0\}$. By Proposition 2.2 they are Cayley polytopes or equivalent to $2\Delta_3$.

If $\Delta' \cong 2\Delta_3$ then $n = \delta = 2$ and without loss of generality $a = 2e_3$. Then $\Delta'' \cong 2\Delta_3$ or $b_3 = -1$. The first case implies $\deg\left(\frac{P}{2}\right) \leq 1$ and $\frac{P}{2}$ has only unimodular facets, which is not possible for a circuit. In the second case we get $P \hookrightarrow 3\Delta_3$.

So let in the following Δ' and Δ'' both be Cayley polytopes.

Case 1: Δ' and Δ'' are pyramids over $P \cap X_3^\perp$.

Here we get $a_3 = -b_3 = 1$. So we may assume $a = e_3$. If $\delta = n = 2$, then $b = e_1 + e_2 - e_3$, $b = 2e_1 + e_2 - e_3$ or $b = e_1 + 2e_2 - e_3$, i.e. $P \hookrightarrow 3\Delta_3$.

Now let $\delta = 1$. As P is a circuit, we get $b_2 = 1$. Then $P \subset \{0 \leq X_2 \leq 1\}$ is a Cayley polytope.

Case 2: Δ' and Δ'' are pyramids.

Let Δ' be a pyramid over $\text{conv}(0, ne_1, a)$. Then $\delta = 1$ and $\text{conv}(0, ne_1, a)$ is a Lawrence polytope. Thus is Δ' a Lawrence polytope and this case is reduced to case 1.

Case 3: Exactly one of the simplices Δ' and Δ'' is a pyramid.

Without loss of generality let Δ'' be a pyramid and Δ' be the Cayley polytope of $\Delta'_1 = \text{conv}(0, ne_1)$ and $\Delta'_2 = \text{conv}(e_2, a)$. Moreover let $0 \leq a_2 < a_3 > 1$. Like in case 1 and 2 we derive $b_3 = -1$, $\delta = 1$ and $\text{Vol}(\Delta') = na_3$.

Also $\text{Vol}(\text{conv}(0, ne_1, a - e_2)) = \text{Vol}(\Delta')$ and hence $\gcd(na_3, n(a_2 - 1)) = na_3$, which implies $a_3 | (a_2 - 1)$. So $a_2 = 1$. As P is a circuit, we conclude $b_2 = 0$. Hence $P \subset \{0 \leq X_2 \leq 1\}$ is a Cayley polytope.

Case 4: Neither Δ' nor Δ'' is a pyramid.

Like in case 3 we may assume $a_2 = 1$ and $b_3 | (b_2 - 1)$. Consider the projection of P to $X_1^\perp$. As Δ' is not a pyramid, we notice $a_3 \geq 2$. Then we get $b_2 \leq 0$ and $b_3 - a_3 b_2 < 0$, because P is a circuit. Now $-b_3 | (1 - b_2)$, which is smaller than $1 + \frac{-b_3}{a_3}$. This implies $b_3 = -1$, which is a contradiction. $\square$

PROPOSITION 3.3. Let $P \subset \mathbb{R}^3$ be a white lattice polytope. Then is P a Cayley polytope or $\text{Vol}(P) \leq C^{(3,3)}$.

PROOF. If $2\Delta_2 \hookrightarrow P$, then by Proposition 2.2 every lattice point of P will have at most distance 2 from $2\Delta_2$. This bounds P . Thus let $2\Delta_2 \not\hookrightarrow P$. Then will every facet of P be a Lawrence polytope.

Let $\text{conv}(0, ne_3)$, $n \in \mathbb{N}$ be the longest edge of P , with adjacent facets F_1, F_2 . By Howe's Theorem 1.2 and the Propositions 2.2, 3.2 we may suppose $n \geq 2$ and

that P has more than 5 vertices. Choose some vertices $a \in F_1$, $b \in F_2$ such that $\Delta := \text{conv}(0, a, b, ne_3)$ is a 3-dimensional lattice simplex.

If there is no lattice point $p \in P \setminus \{F_1 \cup F_2\} \cap \mathbb{Z}^3$, then P contains a pyramid and will also be a Cayley polytope by Proposition 3.1. So let $p \in P \setminus \{F_1 \cup F_2\} \cap \mathbb{Z}^3$.

By Proposition 3.2 we see that $\text{conv}(0, ne_3, a, b, p)$ is a Cayley polytope. So there are possibilities:

- A The distance of p and b to F_1 is 1.
- B The distance of p and a to F_2 is 1.
- C A plane through a, b and p has distance 1 to 0 and ne_3 .

If there is a lattice point $q \in P \cap \mathbb{Z}^3$ of type A, not of type B and a lattice point $q' \in P \cap \mathbb{Z}^3$ of type B, not of type A, then we may assume $a = e_1$ and $b = e_2$. But then is $\text{conv}(0, ne_3, q, q')$ not a Cayley polytope, which is a contradiction to Proposition 2.2.

If there is a lattice point $q \in P \cap \mathbb{Z}^3$ of type A or type B and a lattice point $q' \in P \cap \mathbb{Z}^3$ of type C, then we may assume $a = e_1$ and $b = e_2$. But then there is no possible position for q' .

So it is only possible to have further lattice points $p \in P \setminus \{F_1 \cup F_2\} \cap \mathbb{Z}^3$ of either type A or type B or type C. This means that P is a Cayley polytope. $\square$

4. Lattice Polytopes Without Interior Lattice Points

In the following it remains to consider lattice polytopes having at least one facet with an interior lattice point in order to prove Theorem 1.3.

REMARK 4.1. Let $F \subset \mathbb{R}^2$ be a lattice polygon with $\text{Vol}(F) \leq 4$ and $F^\circ \cap \mathbb{Z}^2 \neq \emptyset$. Pick's formula states $\text{Vol}(F) = |F \cap \mathbb{Z}^2| + |F^\circ \cap \mathbb{Z}^2| - 2$. So we derive $|F^\circ \cap \mathbb{Z}^2| = 1$ and F is one of the following:

$$\{\text{conv}(e_1, e_2, -e_1 - e_2), \text{conv}(\pm e_1, 2e_2), \text{conv}(\pm e_1, \pm e_2), \text{conv}(e_1, e_2, \pm(e_1 + e_2))\}$$




FIGURE 3. The 4 smallest lattice polygons with an interior lattice point

PROPOSITION 4.2. Let $P = \text{conv}(F, d) \subset \mathbb{R}^3$ be a lattice simplex without interior lattice points and $|F^\circ \cap \mathbb{Z}^3| = 1$, $d \in \mathbb{Z}^3$, $d_3 > 0$. Let every further facet of P be without interior lattice points. Then d_3 is bounded by a constant $C^{(4.2)} > 0$.

REMARK 4.3. In fact, it is possible to show that $C^{(4.2)} = 6$.

PROOF. There is a lattice point $p \in \mathbb{Z}^3 \setminus P$ such that $\text{conv}(p, P) = P \cup \text{conv}(p, F)$ and $|\text{conv}(p, P)^\circ \cap \mathbb{Z}^3| = 1$. From Theorem 1.1 it follows that d_3 is bounded. $\square$

It is easy to generalize this idea in order to prove the following:

PROPOSITION 4.4. Let $F \subset \mathbb{R}^n$ be an $(n-1)$ -dimensional lattice polytope with $i > 0$ interior lattice points and $d \in \mathbb{Z}^n$ such that $P = \text{conv}(F, d)$ is an n -dimensional lattice polytope without interior lattice points. Then the volume of P is bounded by a constant depending only on n and i .

PROPOSITION 4.5. Let $P \subset \mathbb{R}^3$ be a lattice polytope and $F \subset P$ be a facet of P with $|F^\circ \cap \mathbb{Z}^3| > 0$. Then P is a Cayley polytope, P can be projected to the double unimodular 2-simplex or $\text{Vol}(P) < C^{(4.5)}$.

PROOF. Let without loss of generality $F = P \cap X_3^\perp$ and $X_3|_P \geq 0$. Define $s := P \cap (X_3 - 1)^\perp$. Assume P is not contained in $\{0 \leq X_3 \leq 1\}$, i.e. $s^\circ \subset P^\circ$.

There is a projection $\pi : \mathbb{Z}^3 \rightarrow \mathbb{Z}^2$ mapping s onto an interval I of length $d = \text{vol}(I) \geq 1$ such that $d \leq f := \max_{x,y \in s} |\langle v, x - y \rangle|$, whereas v is a primitive normal vector of I .

If $d = 1$ we will derive $\text{Vol}(P)$ is bounded or $\pi(P) \cong 2\Delta_2$, because there is a lattice point $p \in P \cap \mathbb{Z}^3$ satisfying $p_3 > 1$ and $\pi(P)$ is a lattice polytope. So let $d > 1$. By $d \leq f$ and $s^\circ \cap \mathbb{Z}^3 = \emptyset$ we receive after some computation $d \leq 2$. We will distinguish the following three cases:

Case 1: $1 < d \leq \frac{3}{2}$.

As $\pi(P)$ is a lattice polytope and there is a lattice point $p \in P \cap \mathbb{Z}^3$ with $p_3 > 1$, we see $p_3 \leq 4 \forall p \in P \cap \mathbb{Z}^3$ and $d \in \{\frac{4}{3}, \frac{3}{2}\}$. By calculating we receive hence $\text{Vol}(s) \leq \frac{16}{3}$. Therefore P is contained in the truncated cone spanned by F and s and contained in $\{0 \leq X_3 \leq 4\}$. This bounds $\text{Vol}(P)$.

Case 2: $\frac{3}{2} < d \leq 2$, $\text{Vol}(F) > 4$.

By computation we receive $\text{Vol}(s) \leq \frac{9}{2}$. As $\text{Vol}(F) \geq 5$ we conclude $p_3 < 20 \forall p \in P \cap \mathbb{Z}^3$. Therefore P is contained in the truncated cone spanned by F and s and contained in $\{0 \leq X_3 \leq 19\}$. This bounds $\text{Vol}(P)$.

Case 3: $\frac{3}{2} < d \leq 2$, $\text{Vol}(F) \leq 4$.

Again $\text{Vol}(s) \leq \frac{9}{2}$. By Remark 4.1, F is equivalent to one of the 4 polygons from Remark 4.1. Moreover we may assume that every facet of P with interior lattice points is one of them. Let $p \in P \cap \mathbb{Z}^3$ with p_3 maximal and consider the lattice polytope $\Delta_p := \text{conv}(F, p)$.

If F is the only facet of Δ_p with an interior lattice point, we will derive $p_3 \leq 6$ by Remark 4.3.

Else let there be a facet $F' \neq F$ of Δ_p with an interior lattice point $q \in F'^\circ \cap \mathbb{Z}^3$. As P has no interior lattice points, we derive $F' \subset \partial P$. Thus F' is one of the four lattice polygons from Remark 4.1. As $q_3 \leq 6$ again, we receive $p_3 \leq 3q_3 = 18$.

Therefore P is contained in the truncated cone spanned by F and s and contained in $\{0 \leq X_3 \leq 18\}$. This bounds $\text{Vol}(P)$. $\square$

PROOF OF THEOREM 1.3. If a lattice polytope $P \subset \mathbb{R}^3$ without interior lattice points is neither a Cayley polytope nor can be projected to the double unimodular 2-simplex, it will follow from the Propositions 3.3 and 4.5 that $\text{Vol}(P) \leq \max\{C^{(3,3)}, C^{(4,5)}\}$. We conclude that there is only a finite number of such exceptional polytopes by Theorem 1.1. $\square$

References

- [1] J.R. Arkinstall, *Minimal requirements for Minkowski's theorem in the plane. I, II*, Bull. Austral. Math. Soc. **22** (1980), no. 2, 259–283. MR 598699.
- [2] V. Batyrev, B. Nill, *Multiples of lattice polytopes without interior lattice points*, Moscow Math. J. **7** (2007), 195–207.
- [3] M. Franz, *Convex – a Maple package for convex geometry*, version 1.1 (2006), available at <http://www-fourier.ujf-grenoble.fr/~franz/convex/>
- [4] C. Haase, G.M. Ziegler, *On the maximal width of empty lattice simplices*, Europ. J. Comb. **21**(2000), no 1, 111–119.
- [5] D. Hensley, *Lattice vertex polytopes with interior lattice points*, Pacific J. Math **105** (1983), no. 1, 183–191.
- [6] A.G. Khovanskii, *Newton polyhedra and the genus of complete intersections*, Funct. Anal. Appl. **12** (1978), no. 1, 38–46.
- [7] A.G. Khovanskii, *Newton polygons, curves on torus surfaces, and the converse Weil theorem*, Russian Math. Surveys **52** (1997), 1251–1279.
- [8] R.J. Koelman, *The number of moduli of families of curves on toric surfaces*, 1991, Proefschrift, Katholieke Universiteit te Nijmegen.
- [9] J.C. Lagarias, G.M. Ziegler, *Bounds for lattice polytopes containing a fixed number of interior points in a sublattice*, Canad. J. Math **43** (1991), no. 5, 1022–1035.
- [10] D.R. Morrison, G. Stevens, *Terminal quotient singularities in dimension three and four*, Proc. Amer. Math. Soc. **90** (1984), no. 1, 15–17.
- [11] B. Reznick, *Clean lattice tetrahedra*, Preprint, math.CO/0606227v1.
- [12] A. Sebő, *An introduction to empty lattice-simplices*, Preprint, November 1998.
- [13] H.E. Scarf, *Integral polyhedra in three space*, Math. Oper. Res. **10** (1985), 403–438.
- [14] J. Schicho, *Simplification of surface parametrizations – a lattice polygon approach*, J. Symb. Comp. **36**, (2003), 535–554.
- [15] G.K. White, *Lattice tetrahedra*, Canad. J. Math. **16** (1964), 389–396.

DEPARTMENT OF MATHEMATICS AND PHYSICS, UNIVERSITY OF TÜBINGEN, AUF DER MORGENSTELLE 10, D-72076 TÜBINGEN, GERMANY

E-mail address: jaron@mail.mathematik.uni-tuebingen.de